










## **Shining X-rays on Catalytic Reactors: Importance of Time and Spatially Resolved Studies**

Jan-Dierk Grunwaldt

Institute for Chemical Technology and Polymer Chemistry, Campus South







KIT – University of the State of Baden-Wuerttemberg an

www.kit.edu

#### Overview





- Introduction: Setting the scene
- Time resolved studies
- Scanning and full-field X-ray microscopy in the µm and mm regime
- Spatiotemporal studies
- Summary and outlook

2 28.06.2010

Shining X-rays on catalytic reactors | Grunwaldt










#### Imaging working catalysts



- Improvement of catalysts triggered by the microscale
- Equally important is the "macroscale"
  - Real catalysts are in large scale fixed bed reactors
    - Pressure drop needs to be minimized
    - Strength of catalyst pellets
    - Optimized diffusion parameters







A.T. Bell, Science 299, 1688 (2003)

(Haldor Topsøe)

28.06.2010

Shining X-rays on catalytic reactors | Grunwaldt


Institute for Chemical Technology and Polymer Chemistry

### **Imaging working catalysts**




- Improvement of catalysts triggered by the microscale
- Equally important is the "macroscale"
  - Real catalysts are in large scale fixed bed reactors
    - Pressure drop needs to be minimized
    - Strength of catalyst pellets
    - Optimized diffusion parameters
  - Temperature and concentration gradients in catalytic reactors

# Partial oxidation of methane over 2.5%Rh/Al<sub>2</sub>O<sub>3</sub>




Partial oxidation of methane over Pd



28.06.2010 Shining X-rays on catalytic reactors | Grunwaldt



### Imaging working catalysts - time resolution Long-term dynamic effects Sintering and ageing Restructuring Formation of BaCeO<sub>3</sub>, Pt<sub>x</sub>Ba<sub>1-x</sub>CeO<sub>3</sub> in NSR-catalysts — Pt/Ba/CeO, raw material .... calcined at 800°C for 1h ... calcined at 1000°C for 8h after calcination to 1200°C BaCeO. Progress of formation / % FT (k3-weighted) 0.015 BaAI,O 0.010 0.005 M. Casapu, J.-D. Grunwaldt, et al., Top. Catal. 42, 3 (2007), J. Catal, 251, 48 (2007) 28.06.2010 Shining X-rays on catalytic reactors | Grunwaldt Institute for Chemical Technology and Polymer Chemistry






#### Imaging working catalysts - time resolution




- Long-term dynamic effects
  - Sintering and ageing, restructuring
- Rapid structural changes
  - Temperature programmed reaction
  - Ignition, extinction
  - Oscillating reactions
  - Activation of catalysts (e.g. reduction)

Total oxidation of %/methane over 10%Pd/ZrO<sub>2</sub>

28.06.2010



Grunwaldt, et al. Chem. Commun., 4635 (2007).
Shining X-rays on catalytic reactors | Grunwaldt




Grunwaldt and Baiker, Catal. Lett. 99, 5 (2005).


Institute for Chemical Technology and Polymer Chemistry

## Shining X-rays on catalysts – best compromise?





$$\tau_{film} \cong \frac{\rho_{CuO} \cdot \frac{d}{2}}{k_{m} \cdot c_{H2}} \longrightarrow 6.5 \text{ s}$$



$$\tau_{film} \cong \frac{\rho_{CuO} \cdot R_p}{3k_m \cdot c_{H2}} \longrightarrow 1.7 \text{ ms}$$

D<sub>e</sub>=10<sup>-6</sup> to 10<sup>-8</sup> m<sup>2</sup>/s

$$\tau_{pore} \cong \frac{\rho_{CuO} \cdot (\frac{\alpha}{2})^2}{2D_e \cdot c_{H2}} \longrightarrow \frac{1}{2D_e \cdot c_{H2}}$$

$$\tau_{pore} \cong \frac{\rho_{\textit{CuO}} \cdot R_p^2}{6D_e \cdot c_{H2}} \longrightarrow \begin{array}{c} \textbf{0.02 to 2 s} \\ \text{Grunwaldt et al., PCCP 6, 3037 ( 2004).} \end{array}$$

**12** 28.06.2010

Shining X-rays on catalytic reactors | Grunwaldt



#### Catalytic partial oxidation of methane



### $CH_4 + \frac{1}{2}O_2 \rightarrow CO + 2H_2$

#### Active Catalysts: Supported noble metals, Ni-catalysts [1]

- Alternative route to steam/ autothermal reforming [2]
- Synthesis gas for production of liquid fuels
- Useful in stationary SOFCs

[1] A.P.E. York et al., Topics Catal. 22, 345 (2003)
Hickmann and Schmidt, Science 259, 343 (1993)
[2] J.R. Rostrup-Nielsen, J. Sehested, J.K. Norskov, Adv. Catal. 47, 65 (2002)

Flaring of natural gas



From: L. Plass, S. Reimelt, CIT 79, p. 561 (2007)

1 101111 21 1 1000; C. 1101111011; C. 1 1 0; p. 00 1 (2001)

**14** 28.06.2010


Shining X-rays on catalytic reactors | Grunwaldt

Institute for Chemical Technology and Polymer Chemistry

## Catalytic partial oxidation of methane: Ignition of the reaction



### $CH_4 + \frac{1}{2}O_2 \rightarrow CO + 2H_2$




2.5%Rh-2.5%Pt/Al<sub>2</sub>O<sub>3</sub>, prepared by flame synthesis 6% CH<sub>4</sub>, 3% O<sub>2</sub> in He, ramp rate: 5 K/min


Grunwaldt and Baiker, Catal. Lett. 99, 5 (2005).

**15** 28.06.2010


Shining X-rays on catalytic reactors | Grunwaldt

















































































































## Going to the third dimension: X-ray tomography on shaped catalysts





3D-reconstruction of 0.5%Pd/Al<sub>2</sub>O<sub>3</sub>, shell-impregnated catalyst pellet of cylindrical shape; sample was stuck from one side on the sample holder, field of view: 3.54 mm x 2.36 mm; total reconstructed volume of 3.54 x 3.54 x 2.36 mm<sup>3</sup>.

44


28.06.2010


Shining X-rays on catalytic reactors | Grunwaldt

Institute for Chemical Technology and Polymer Chemistry

## Going to the third dimension: X-ray tomography on shaped catalysts







3D-reconstruction of Cu/Al<sub>2</sub>O<sub>3</sub>, (a) 1 min and (b) 10 min with CuCl<sub>2</sub>-solution impregnated catalyst pellet; field of view: 3.54 mm x 2.36 mm.


J.-D. Grunwaldt, et al., Catal. Today 145, 267 (2009).


45

28.06.2010


Shining X-rays on catalytic reactors | Grunwaldt
















































#### **Summary and conclusions**



- Dynamic & in situ studies are important for structural studies both on a micro- and nanoscale
- Hard X-ray based techniques advantageous for gas phase, high temperature, liquid phase and elevated pressure
- We need in future
  - Full EXAFS spectra
  - X-ray absorption tomography under in situ conditions
  - Better spatial resolution
  - Combination of full-field and scanning X-ray microscopy
  - Combination of electron and X-ray microscopy
  - Time-resolution with spatial resolution

28.06.2010

Shining X-rays on catalytic reactors | Grunwaldt

Institute for Chemical Technology and Polymer Chemistry

### **Acknowledgements**









- KIT: The new group especially Matthias Bauer and Alexey Boubnov
- DTU-KT: Matthias Beier, Martin Høj, David Mogensen, Anna Filipu, Bjørn Maribo, Jon Christensen
- •CEN: Rafal Dunin-Borkowski, Thomas Hansen, Jacob Wagner



- Department of Chemistry and Applied Biosciences: Alfons Baiker, Matteo Caravati, Bertram Kimmerle, Stefan Hannemann, and many more
- Mechanical and electronic workshop



• Universität Wuppertal: Prof. R. Frahm, Jan Stötzel, Dirk Lützenkirchen-Hecht



• RWTH Aachen/TU Dresden: Prof. Dr. C.G. Schroer, Prof. B. Lengeler



- PERGAM-SUISSE AG: Infrared-Thermography
- Financial Support: EU-projects, especially R113-CT-2004-506008, DANSCATT, Technical University of Denmark, Danish Research Council, Haldor Topsøe, ETH Zürich, Karl-Winnacker-Fond

28.06.2010

Shining X-rays on catalytic reactors | Grunwaldt



